

# General compatibility of nucleoside triphosphates with rolling circle amplification

Marcel Hollenstein\*


Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, CH-3012 Bern, Switzerland.

[hollenstein@deb.unibe.ch](mailto:hollenstein@deb.unibe.ch)

## Introduction

- Besides PCR, rolling circle amplification (RCA) is the most used amplification method of DNA. In RCA, single-stranded DNA nanocircles serve as virtually infinite templates for short primers, generating long, ssDNA products containing multiple copies of the reverse complement of the template.<sup>1,2</sup> RCA has found a variety of applications including sensing, diagnostics, and nanotechnology.<sup>2</sup>
- Modified nucleoside triphosphates (dN\*TPs) represent a versatile vector for the introduction of chemical diversity into nucleic acids.<sup>3</sup> Surprisingly, while dN\*TPs have been engaged in a large palette of applications including the selection of aptamers<sup>4</sup> and electrochemical tagging of nucleic acids,<sup>5</sup> only few examples combining dN\*TPs and RCA have been reported.<sup>6</sup>
- Herein, the compatibility of dN\*TPs modified at any position of the scaffold with rolling circle amplification method was assessed. This method was used to generate long, nuclease-resistant and fully-modified cytosine-rich mimics of telomeric DNA.<sup>7</sup>

## Design of the method



**Figure 1.** A) Chemical structures of the modified nucleoside triphosphates used in rolling circle amplification; B) Schematic representation of RCA with dN\*TPs: the linear template precursors **T1-3** are circularized with CircLigase™. The nanocircles **cT1-3** served as templates for RCA, along with a suitable primer, dN\*TPs, and a DNA polymerase.


## Polymerase screen

Various DNA polymerases (A- and B-family) were screened to assess whether they could extend the primer in the presence of one or multiple dN\*TPs under RCA conditions. Only the 9°N<sub>m</sub> DNA polymerase is capable of extending the primer in the presence of *all* modifications with high yields.

The 9°N<sub>m</sub> DNA polymerase was also shown to be compatible with a broad variety of templates (Figure 2) and could be used to generate fully modified, long, cytosine-rich mimics of telomeric DNA.

Primer extension reactions with linear templates and a Sanger-sequencing experiment clearly demonstrated that the products emanate from a rolling circle mechanism.


## Generation of fully modified RCA products



**Figure 2.** Analysis (PAGE 5%) of RCA products obtained with templates **cT1-3** and the 9°N<sub>m</sub> DNA polymerase. Lane 1: α-thio dNTPs **6**; lane 2: 2'-fluoro-rNTPs **5**; lane 3: base-modified dN\*TPs **1-3**; lane 4: natural dNTPs.

## Serum resistance

The modified RCA products all display a higher stability in FBS than the corresponding unmodified products:



**Figure 3.** Gel image (PAGE 5%) showing the stability of RCA products in 10% FBS (37°C). Line 1 natural dNTPs; line 2: dN\*TPs **1-3**; line 3: α-thio-dNTPs **6**; line 4: 2'-fluoro-rNTPs **5**. A: template **cT1**; B: template **cT3**.

## Conclusions

- dN\*TPs are compatible with RCA, regardless of the location and nature of the modification and the topology of the template.
- The combination of RCA and dN\*TPs yields long, single-stranded and modified products that have an increased stability in serum.
- This method could be used to develop new biosensors, biocatalysts based on DNazymes, and nanomaterials.

## References

- M. M. Ali, F. Li, Z. Zhang, K. Zhang, D.-K. Kang, J. A. Ankrum, X. C. Le, W. Zhao, *Chem. Soc. Rev.* **2014**, 43, 3324–3341.
- W. Zhao, M. M. Ali, M. A. Brook and Y. Li, *Angew. Chem., Int. Ed.* **2008**, 47, 6330–6337
- a) M. Hocek, *J. Org. Chem.* **2014**, 79, 9914–9921; b) M. Hollenstein, *Molecules* **2012**, 17, 13569–13591.
- S. Diafa, M. Hollenstein, *Molecules* **2015**, 20, 16643–16671.
- M. Hocek, M. Fojta, *Chem. Soc. Rev.* **2011**, 40, 5802–5814.
- A. Baccaro, A. Marx, *Chem. Eur. J.* **2010**, 16, 218–226
- M. Hollenstein, *Org. Biomol. Chem.* **2015**, 13, 9820–9824